arXiv:1211.4737v1 [cond-mat.mes-hall] 20 Nov 2012

Thermodynamic and quantum bounds on nonlinear thermoelectric devices

Robert S. Whitney[1]

[1] *Laboratoire de Physique et Modélisation des Milieux Condensés (UMR 5493), Université Grenoble 1 and CNRS, Maison des Magistères, BP 166, 38042 Grenoble, France.*

(Dated: November 21, 2012)

I consider the non-equilibrium transport of electrons through a quantum system with a thermoelectric response. This system may be any nanostructure or molecule modeled by the nonlinear scattering theory which includes Hartree-like electrostatic interactions exactly and certain dynamic interaction effects (decoherence and relaxation) phenomenologically. I derive three fundamental bounds for such quantum systems coupled to multiple macroscopic reservoirs, one of which may be superconducting. These bounds affect nonlinear heating effects (such as Joule heating), and work and entropy production. Two bounds correspond to the first law and second law of thermodynamics in classical physics. The third bound is *quantum* (wavelength dependent), and is as important as the thermodynamic ones in limiting the capabilities of mesoscopic heat-engines and refrigerators. The quantum bound also leads to Nernst's unattainability principle that the quantum system cannot cool a reservoir to absolute zero in a finite time, although it can get exponentially close.

PACS numbers: 73.63.-b, 05.70.Ln, 72.15.Jf, 84.60.Rb

I. INTRODUCTION

In classical physics, currents in resistive circuits are dissipative; they radiate Joule heat which increases the entropy of the environment of the circuit. Devices with thermoelectric responses[1–3] can be used as heat-engines (using heat-flows to create electrical power) or heat pumps (using electrical power to create heat-flows). These too are dissipative, since they increase the entropy of the environment (at least a little bit) during their operation. The laws of thermodynamics give us strict bounds on the capabilities of such thermoelectric devices.

There is currently great interest in using quantum systems as thermoelectric heat-engines or refrigerator, particularly nanostrutures[4,6–13] and molecules[14–19]. However quantum responses are often non-local, since thermal equilibration occurs on lengthscales very much bigger than the size of the system, with equilibrium only being established deep in the reservoirs coupled to the system. Classical models of transport account neither for this non-locality, nor for the wave-nature of quantum particles. It is natural to ask what the bounds are for these quantum devices, and how they compare to the bounds on classical devices. There are many works asking whether quantum systems are bounded by the laws of thermodynamics[20–27]; particularly for systems modeled by Markovian or Bloch-Redfield master equations, see Refs. [20,28–30] and references therein. However the fundamental bounds on *thermoelectric* quantum systems remains an open question[27]. Here, I consider the fully nonlinear response of a broad class of thermoelectric quantum systems, and show that they obey the bounds imposed by the laws of thermodynamics, while also suffering a quantum bound on their behavior.



Figure 1: Quantum systems with $K_{\rm n} = 3$ normal reservoirs and one superconducting (sc) reservoir. Quantum systems may be (a) nanostructures such as those defined by electrostatic gates on top of a two-dimensional electron gas (2deg), or (b) molecules. Charge and heat current, I_i and J_i, are positive if they flow from the quantum system to reservoir i.

The Landauer-Büttiker scattering theory[33,34] is a widely-used theory for modeling such electron flows through quantum systems; giving DC and AC conductances[35,36], and nonlinear effects[37–40]. It has equally been used to get linear thermal and thermoelectric effects[41–46]. However it is a delicate question whether it captures dissipation and the associated laws of thermodynamics (or their quantum equivalent). The standard answer is that the theory accurately models *reversible* scattering processes, while assuming that equilibriation occurs (irreversibly) in the reservoirs via dissipative processes which the theory does not detail. Hence, one might suspect that the theory would not capture how dissipation produces entropy. This suspicion is wrong, the nonlinear scattering theory is sufficient to give information about entropy production, despite the simplistic way in which it treats the physics in the reservoirs. Indeed, Ref. [27] showed the second law of thermodynamics to be a remarkably direct outcome of a nonlinear "DC" scatter-

ing theory[31] for any quantum system of *non-interacting* electrons between two reservoirs[32]. Energy conservation leds to the systems also obeying the first law.

Here I show that the proof in Ref. [27] works equally well for a non-equilibrium scattering theory which includes Hartree-like *electron-electron interactions* (electrostatic charging effects) in a self-consistent and gauge-invariant manner. This theory is the analogue for heat-currents[13] of the nonlinear scattering theory for charge-currents[37–40]; although heat-currents are not conserved when charge-currents are. Such electrostatic interactions cannot be ignored for the nonlinear response[37–40], and entropy production is a purely nonlinear effect (vanishing in the linear-response regime). By keeping only electrostatic interaction effects, this approach is by no means exact, however it is a reasonable approximation for any system where the single-electron charging energy is small compared to temperature or broadening of the quantum system's levels due to the coupling to the reservoirs[37]. Within this approach, I prove that the DC response of any quantum system coupled to an arbitrary number of reservoirs — one of which may be superconducting — obeys the laws of thermodynamics.

The extension to many reservoirs (from two reservoirs in Ref. [27]) requires no new physics, but is technically complicated, because one can no longer use the simple form of two-terminal Onsager relations. However it extends our physical understanding of thermodynamic laws in quantum-transport in two important ways.

(a) Using the generalization to many reservoirs, one can include dynamics interaction effects (decoherence and relaxation) phenomenologically as additional fictitious reservoirs[25,47,48]. Performing such an analysis, I conclude that the classical laws of thermodynamics hold for partially coherent (as well as fully coherent) quantum-transport.

(b) A superconducting (sc) reservoir has no analogue in classical mechanics, because the condensate is a macroscopic quantum state. In addition, charge-current can flow into (or out of) this condensate, but heat-current cannot. Thus a charge-current can pass into the sc reservoir, $I_{\rm sc}$ without any change in that reservoir's entropy; $\dot{S}_{\rm sc} = J_{\rm sc}/T_{\rm sc} = 0$. Despite this, I show that the laws of thermodynamics hold in the presence of such a reservoir.

I also derive a *quantum bound* (qb) on the thermoelectric response. It tells us that no matter how much DC power one supplied to any device, it cannot extract heat from the ith reservoir faster than the bound $J_i^{\rm qb} \propto N_i \times (k_{\rm B}T_i)^2$, where T_i is the reservoir temperature, and the contact between the quantum device and the reservoir has N_i transverse modes. This bound is "quantum"; i.e. it is only relevant when $N_i^{-1} \sim \lambda_{\rm F}^2/A_i \neq 0$, for electron wavelength $\lambda_{\rm F}$, and contact cross-section A_i. The quantity $J_i^{\rm qb}$ corresponds to the heat-current out of the ith reservoir when it is coupled to a zero-temperature reservoir with the same chemical-potential, through a contact containing N_i modes. In such a context, it is a fermionic analogue[49] of the Stefan-Boltzmann law of black-body radiation, for which $J \propto T^4$.

For thermoelectric refrigerators, the quantum and thermodynamic bounds *compete*, and the quantum bound can be a stronger constraint than that given by the thermodynamic bounds (the Carnot limit). In contrast, the bounds *combine* to give an upper limit on the power that a thermoelectric heat-engine can produce.

I find that the quantum bound means that the systems obey Nernst's *unattainability principle* (one of the two statements forming the third law of thermodynamics). This principle claims that a reservoir cannot be refrigerated to absolute zero in a finite time, see for example Refs. [21,29,30] for physical and Refs. [50,51] for mathematical overviews. The quantum bound means that the systems considered here have (at best) the critical behavior; i.e. the reservoir *cannot* achieve zero temperature in a finite time, but it can get exponentially close. This rules out such systems as candidates for violating the unattainability principle in the manner Ref. [30] proposes.

II. EXPLICIT FORM OF THE BOUNDS

In this section I give the three bounds, before deriving them latter in the article. I consider an arbitrary quantum system coupled to any number of reservoirs, one of which may be superconducting. Normal reservoirs are treated as free-electron gases in thermal equilibrium, while a superconducting reservoir is treated as a condensate of Cooper pairs[52]. The heat-current into reservoir i is J_i. The electrical power flowing into reservoir i is $P_i = V_i I_i$, with V_i being the bias on the reservoir, and I_i being the charge-current flowing into it. The rate of change of entropy in the ith reservoir is defined as $\dot{S}_i = J_i/T_i$ for reservoir temperature T_i.

I consider the DC response of the quantum system[31], and assume that any variations of the energy or entropy of the quantum system over time are small enough to ignore. Defining, $\dot{E}_{\rm total}$ and $\dot{S}_{\rm total}$ as the rate of total energy and entropy flow *out* of the quantum system into the reservoirs, averaged over long times. I show that scattering theory reproduces the thermodynamic laws,

$$\dot{E}_{\rm total} \;=\; \sum_i (J_i + P_i) \;=\; 0 \qquad [1^{\rm st}\ \text{law}] \qquad (1)$$

$$\dot{S}_{\rm total} \;=\; \sum_i J_i/T_i \;\geq\; 0 \qquad [2^{\rm nd}\ \text{law}] \qquad (2)$$

where the sum is over all reservoirs. The quantum bound (qb) on the heat current into reservoir i is

$$J_i \geq -J_i^{\rm qb} \equiv -\frac{\pi^2}{6h} N_i \,(k_{\rm B}T_i)^2 \qquad [\text{quantum}] \qquad (3)$$

where the quantum system couples to reservoir i through N_i transverse modes, and this equation defines $J_i^{\rm qb}$. Sections IX and X discuss how these bounds act as limits on heat-engines and refrigerators.

III. NONLINEAR SCATTERING THEORY FOR HEAT TRANSPORT

Here I consider the nonlinear version of the Landauer-Büttiker scattering theory of transport, which includes (in a gauge-invariant and self-consistent manner) those Hartree-type electron-electron interaction effects which can be considered as electrostatic charging effects. It has been used for AC charge-transport[35,36] and nonlinear DC charge-transport[37–40]. Here, as in Ref. [13], I apply it to nonlinear DC heat-transport.

One starts[37–40] with the scattering matrix, $\mathcal{S}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right)$, of a ν-particle entering the quantum system from reservoir j to a μ-particle leaving via reservoir i, where μ and ν can be either electron (with charge $e^{\text{-}}$) or hole (with charge $-e^{\text{-}}$). The superconducting (sc) reservoir is modeled by Andreev reflection of electrons into holes and vice versa. Here an electron with energy ϵ is an *occupied* conduction-band state at energy $\epsilon + e^{\text{-}}V_{\rm sc}$, while a hole at energy ϵ is an *empty* conduction-band state at energy $-\epsilon + e^{\text{-}}V_{\rm sc}$, where $e^{\text{-}}V_{\rm sc}$ is the superconductor's chemical-potential. Thus these "holes" are not those in Ref. [13], which are in a semiconductor's valence band.

One assumes that the lead to reservoir i has N_i modes independent of ϵ. The number of *open* modes in this lead can still be ϵ-dependent; if a given mode n_i is closed off at a certain ϵ, then it appears as purely reflected in the scattering matrix. If a superconducting reservoir is present, one measures the energy ϵ from the superconductor's chemical-potential $e^{\text{-}}V_{\rm sc}$; otherwise one can take $\epsilon = 0$ however is convenient. I define $\mathbf{v}_{\rm env}$ as the set of all voltages and temperatures in the environment of the quantum system, $\{V_k, T_k\}$, where k runs over all reservoirs and gates. In the nonlinear regime, $\mathcal{S}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right)$ must include the charging effects self-consistently, because it is a function of the charge-distribution in the quantum system, which is in turn a function of $\mathcal{S}$ as well as the set of all voltages and temperatures in $\mathbf{v}_{\rm env}$[37,40].

All the transport properties (both charge and heat currents), are then given by the transmission matrix, $\mathcal{T}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right) = {\rm Tr}\left[\left[\mathcal{S}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right)\right]^\dagger \mathcal{S}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right)\right]$ where the trace is over all transverse modes of the leads connecting the quantum system to the i and j reservoirs. Any self-consistent analysis must of course respect gauge-invariance; i.e. shifting all voltages by the same amount is tantamount to redefining the zero of energy, and thus cannot change the physics. I ensure this by taking ϵ and all reservoir and gate voltages (and any screening potentials which depend on them, as relative to the superconductor's chemical-potential,[53] $e^{\text{-}}V_{\rm sc}$. If there is no superconductor, one can take these relative to another reservoir voltage. In everything that follows, I will assume that $\mathcal{T}^{\mu\nu}_{ij}$ has this gauge-invariant.

There are two properties of $\mathcal{T}^{\mu\nu}_{ij}$, which will be crucial in deriving the results presented in this article. The first property is due to the fact that $\mathcal{T}^{\mu\nu}_{ij}$ is the sum of the modulus squared of the elements of $\mathcal{S}^{\mu\nu}_{ij}$. This property is that for all $i, j, \mu, \nu, \epsilon, \mathbf{v}_{\rm env}$,

$$\mathcal{T}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right) \geq 0. \tag{4}$$

The second property is due to the unitary of $\mathcal{S}^{\mu\nu}_{ij}$ (particle conservation), and is that for all $\epsilon, \mathbf{v}_{\rm env}$,

$$\begin{aligned} 0 &= \sum_{j=1}^{K_{\rm n}} \sum_{\nu} \left[\mathcal{T}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right) - N_i\delta^{\mu\nu}_{ij}\right] \\ &= \sum_{i=1}^{K_{\rm n}} \sum_{\mu} \left[\mathcal{T}^{\mu\nu}_{ij}\left(\epsilon,\mathbf{v}_{\rm env}\right) - N_i\delta^{\mu\nu}_{ij}\right], \end{aligned} \tag{5}$$

where the i or j sums are over the $K_{\rm n}$ normal reservoirs, while the μ or ν sums are over ± 1, with $+1$ for electrons (e) and -1 for holes (h).

The charge current into the ith reservoir counts the charge $\mu e^{\text{-}}$ that enters or leaves that reservoir. The heat current into the ith reservoir counts the excess-energy $(\epsilon - \mu e^{\text{-}}V_i)$ that enters or leaves that reservoir[43]. The charge current, I_i, and the heat-current, J_i, from the quantum system into the ith normal reservoir are

$$I_i = \sum_{j\mu\nu} \int_0^\infty \frac{{\rm d}\epsilon}{h}\, \mu e^{\text{-}} \left[\mathcal{T}^{\mu\nu}_{ij}(\epsilon) - N_i\delta^{\mu\nu}_{ij}\right] f^\nu_j(\epsilon), \tag{6}$$

$$J_i = \sum_{j\mu\nu} \int_0^\infty \frac{{\rm d}\epsilon}{h}\, (\epsilon - \mu e^{\text{-}}V_i) \left[\mathcal{T}^{\mu\nu}_{ij}(\epsilon) - N_i\delta^{\mu\nu}_{ij}\right] f^\nu_j(\epsilon). \tag{7}$$

Here j is summed over the $K_{\rm n}$ normal reservoirs, while μ and ν are summed over e $(+1)$ and h (-1). Here $\delta^{\mu\nu}_{ij}$ is shorthand for the product of Kronecker δ-functions, $\delta_{ij}\delta_{\mu\nu}$, while the Fermi function for ν-particles entering the quantum system from reservoir j, is

$$f^\nu_j(\epsilon) = \left(1 + \exp\left[(\epsilon - \nu e^{\text{-}}V_j)/(k_{\rm B}T_j)\right]\right)^{-1}. \tag{8}$$

If a superconductor is present, then the charge-current into it (in the form of Cooper pairs), $I_{\rm sc}$, is given by Kirchoff's law of current conservation. In contrast, heat current is not conserved. However heat flow through a boundary at the surface of a superconducting reservoir must be zero, since an electron at energy ϵ is Andreev reflected as a hole with the same energy. Thus

$$I_{\rm sc} = -\sum_j I_j, \qquad J_{\rm sc} = 0. \tag{9}$$

For $I_{\rm sc}$ the sum is over the $K_{\rm n}$ normal reservoirs.

Comparing Eqs. (6) and (7), one easily see that

$$J_i = -V_i\, I_i + \sum_{j\mu\nu} \int_0^\infty \frac{{\rm d}\epsilon}{h}\, \epsilon\, \left[\mathcal{T}^{\mu\nu}_{ij}(\epsilon) - N_i\delta^{\mu\nu}_{ij}\right] f^\nu_j(\epsilon), \tag{10}$$

Summing the heat current over all reservoirs (noting that $J_{\rm sc} = 0$), one can use Eq. (5) to show that the second term in Eq. (10) cancels, so $\sum_{i=1}^{K_{\rm n}} J_i = -\sum_{i=1}^{K_{\rm n}} V_i\, I_i$.

Note that I have chosen to measure all voltages relative to the superconductor (if present). If instead, one measured the voltages relative to an external reference, then the total heat-current out of the quantum system is

$$J_{\rm total} = -V_{\rm sc}\, I_{\rm sc} - \sum_{i=1}^{K_{\rm n}} V_i\, I_i. \tag{11}$$

Kirchoff's law in Eq. (9) means that changing all voltages by the same amount does not change the total heat flow, $J_{\rm total}$. Thus $J_{\rm total}$ respects gauge-invariance whenever I_i does, where one recalls that I_i is a self-consistent function of[40] $(V_1, T_1, V_2, T_2, \cdots V_N, T_N)$. Whenever $J_{\rm total}$ is non-zero, heat-currents are not conserved in the quantum system; the quantum system is a heat-source for $J_{\rm total} > 0$, or a heat-sink for $J_{\rm total} < 0$.

IV. ZEROTH LAW OF THERMODYNAMICS

The definition of equilibrium is one of the various statements that together form the zeroth law. It is defined as a state for which one can cut the system into parts in any manner, and one will find *no* charge and heat flow between those parts.

A consequence of this is that if a system in equilibrium consists of multiple reservoirs (evidently in equilibrium with each other) all coupled to each other through a quantum system, then there can not be any charge or heat current through that quantum system. It is trivial to use the scattering theory to show that any quantum system, placed between an arbitrary number of reservoirs all in equilibrium with each other, must obey this consequence of the zeroth law. It is sufficient to note that the scattering matrix $\mathcal{S}$ is unitary, this means that the transmission matrix obeys Eq. (5). Substituting this into Eqs. (6,7), one see that charge and heat currents are zero whenever $T_j = T_0$ and $V_j = V_0$ for all j.

V. FIRST LAW OF THERMODYNAMICS AND JOULE HEATING

Eqs. (11) has the form of a *classical* Joule heating term; i.e. *voltage* × *current*. This is due to the fact that energy conservation ensures that the heat-radiated (or absorbed) is equal and opposite to electrical power supplied (or generated). Thus, one arrives directly at the first law of thermodynamics, Eq. (1). However it is important to note that this is only true upon summing over all reservoirs, there is no particular relation between the power into reservoir i, $P_i = V_i I_i$, and the heat-current into that reservoir, given in Eq. (10).

As a first example, consider a quantum system with two normal reservoirs at the same temperature but different biases, Kirchoff's law means that $I_2 = -I_1$, and hence $P_{\rm total} = (V_1 - V_2)I_1$. Then I_1 has the opposite sign to $V_1 - V_2$, and hence $P_{\rm total} < 0$. This means the total heat current out of the quantum system, $J_{\rm total} = -P_{\rm total}$, is positive and corresponds to Joule heating. Note that if reservoir 1 is superconducting all the heat generated flows into reservoir 2. This means that superconductors can be used to filter out heat flow generated by Joule heating (as well as any other heat flow).



Figure 2: A quantum system with a finite thermoelectric response being used as a heat-engine, providing power to a set of loads. the sc reservoir is superconducting, but "load sc" is not (its simply the load coupled to the sc reservoir).

As a second example, the quantum system in Fig. 2 is assumed to have a finite thermoelectric response. If one reservoir is heated it provides a total electrical power of $P_{\rm total} = \sum_i V_i I_i$ to the loads. Then the total heat flow out of the quantum system equals $-P_{\rm total}$ (meaning more heat flows-in than flows-out) and the first law is satisfied.

VI. SECOND LAW OF THERMODYNAMICS

A. System coupled to two reservoirs

Here I reproduce the calculation in Ref. [27] whose outcome is the second law in two-reservoir systems. The Hartree-type interaction effects can be included without any changes to their method. For two reservoirs, the rate of change of entropy is $\dot{S} = (J_1/T_1 + J_2/T_2)$, one can substitute in Eq. (7) and explicitly write out the sum over $j = 1, 2$. There is no Andreev scattering so $\mathcal{T}_{ij}^{\mu\nu} \propto \delta_{\mu\nu}$, hence when there are only two-reservoirs Eq. (5) gives us the following two-terminal Onsager relation, $\mathcal{T}_{11}^{\mu\mu} = -\mathcal{T}_{21}^{\mu\mu} = \mathcal{T}_{22}^{\mu\mu} = -\mathcal{T}_{12}^{\mu\mu}$. As a result,

$$\frac{\dot{S}_{\rm total}}{k_{\rm B}} = \sum_{\mu} \int_0^{\infty} \frac{{\rm d}\epsilon}{h} \big[\xi_{1\mu} - \xi_{2\mu}\big] \times \mathcal{T}_{12}^{\mu\mu}(\epsilon, \mathbf{v}_{\rm env}) \big[f(\xi_{2\mu}) - f(\xi_{1\mu})\big], \tag{12}$$

where I define $\xi_{i\mu} = (\epsilon - \mu e^{-} V_i)/(k_{\rm B} T_i)$, so $f_i^{\mu}(\epsilon) = f(\xi_{i\mu})$ for $f(\xi) = (1 + \exp[\xi])^{-1}$. Since $f(\xi)$ is a monotonically decaying function of ξ, the product of the two square-brackets in Eq. (12) cannot be negative. Taking this together with Eq. (4), one conclude that the second law, $\dot{S}_{\rm total} \geq 0$, is satisfied for such a two-reservoir system.

B. System coupled to any number of reservoirs

In general for more than two reservoirs, there is no such simple relationship between $\mathcal{T}^{\nu\mu}_{ji}$ and $T^{\mu\nu}_{ij}$ as above. Then the derivation leading to the second law is more involved. Here I provide such a derivation; one that uses only the unitarity of $\mathcal{S}^{\mu\nu}_{ij}$, in the form of Eqs. (4,5). Using Eq. (7), I write $\dot{S}_{\rm total} = (k_{\rm B}/h)\int_0^\infty {\rm d}\epsilon\; Y(\epsilon)$ with

$$Y(\epsilon) \;=\; \sum_{ij\mu\nu} \xi_{i\mu}\,\left[\mathcal{T}^{\mu\nu}_{ij}(\epsilon) - N_i\delta^{\mu\nu}_{ij}\right]\, f(\xi_{j\nu}). \tag{13}$$

where i and j sums are over the $K_{\rm n}$ normal reservoirs. As in Eq. (12), I defined $\xi_{i\mu} = (\epsilon - \mu e^{-}V_i)/(k_{\rm B}T_i)$ and $f(\xi) = (1+\exp[\xi])^{-1}$. I now prove that $Y(\epsilon)\geq 0$ for all ϵ, thereby showing the second law, Eq. (2), is satisfied.

One can re-labeling the reservoirs in Eq. (13), giving separate labels to the electron-state entering a given reservoir, and the hole states entering that same reservoir. Thus I replace the sum $\sum_{i=1}^{K_{\rm n}}\sum_{\mu=\pm 1}$ with $\sum_{\tilde{\imath}=1}^{2K_{\rm n}}$, where $\tilde{\imath}$ is related to i and μ as follows. For each i and μ one calculates $\xi_{i\mu}$, and then order the elements in ascending order of $\xi_{i\mu}$. One then labels these states with superscripts, $\tilde{\imath}$, running from 1 to $2K_{\rm n}$, such that $\xi^{(1)} < \xi^{(2)} < \cdots < \xi^{(2K_{\rm n})}$. I then define

$$\xi^{(\tilde{\imath})} \;=\; \xi^{(0)} + \sum_{\widetilde{n}=1}^{\tilde{\imath}} \Delta\xi^{(\widetilde{n})}, \tag{14}$$

$$f(\xi^{(\tilde{\imath})}) \;=\; f(\xi^{(0)}) + \sum_{\widetilde{m}=1}^{\tilde{\jmath}} \Delta f^{(\widetilde{m})}, \tag{15}$$

where $\Delta\xi^{(\widetilde{n})} = \left[\xi^{(\widetilde{n})} - \xi^{(\widetilde{n}-1)}\right]$ and $\Delta f^{(\widetilde{m})} = \left[f(\xi^{(\widetilde{m})}) - f(\xi^{(\widetilde{m}-1)})\right]$. Here $\xi^{(0)}$ is arbitrary, so I take $\xi^{(0)} < \xi^{(1)}$. Then$\Delta\xi^{(\widetilde{n})}\geq 0$ for all $\widetilde{n}$, and $\Delta f^{(\widetilde{m})}\leq 0$ for all $\widetilde{m}$ (since $f(\xi)$ decays monotonically with ξ).

Taking Eq. (13) and replacing the quadruple sum over i,j,μ,ν with a double sum over $\tilde{\imath},\tilde{\jmath}$, I substitute for $\xi^{(\tilde{\imath})}$ and $f(\xi^{(\tilde{\jmath})})$ using Eqs. (14,15). Then

$$\begin{aligned} Y(\epsilon) \;=\; \sum_{\tilde{\imath},\tilde{\jmath}=1}^{2K_{\rm n}} \Bigg[& \xi^{(0)}\;\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\; f(\xi^{(0)}) \\ & + \sum_{\widetilde{n}=1}^{\tilde{\imath}} \Delta\xi^{(\widetilde{n})}\;\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\; f(\xi^{(0)}) \\ & + \sum_{\widetilde{m}=1}^{\tilde{\jmath}} \xi^{(0)}\;\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\; \Delta f^{(\widetilde{m})} \\ & + \sum_{\widetilde{n}=1}^{\tilde{\imath}}\sum_{\widetilde{m}=1}^{\tilde{\jmath}} \Delta\xi^{(\widetilde{n})}\;\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\; \Delta f^{(\widetilde{m})} \Bigg]. \end{aligned}$$

However given Eq. (5), the first three lines of this result give zero when summed over $\tilde{\imath},\tilde{\jmath}$; leaving only the term containing the sum over both $\widetilde{n}$ and $\widetilde{m}$ to evaluate. To do this, I write $\sum_{\tilde{\imath}=1}^{2K_{\rm n}}\sum_{\widetilde{n}=1}^{\tilde{\imath}} = \sum_{\widetilde{n}=1}^{2K_{\rm n}}\sum_{\tilde{\imath}=\widetilde{n}}^{2K_{\rm n}}$ and $\sum_{\tilde{\jmath}=1}^{2K_{\rm n}}\sum_{\widetilde{m}=1}^{j} = \sum_{\widetilde{m}=1}^{2K_{\rm n}}\sum_{\tilde{\jmath}=\widetilde{m}}^{2K_{\rm n}}$, after which

$$Y(\epsilon) \;=\; \sum_{\widetilde{n},\widetilde{m}=1}^{2K_{\rm n}} \Delta\xi^{(\widetilde{n})}\;\Delta f^{(\widetilde{m})} \sum_{\tilde{\imath}=\widetilde{n}}^{2K_{\rm n}}\sum_{\tilde{\jmath}=\widetilde{m}}^{2K_{\rm n}} \left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right].$$

For $\widetilde{n} > \widetilde{m}$, one can prove that the sum over $\tilde{\jmath}$ is negative (or zero) for all $\tilde{\imath}$. The proof requires noting that every sum over $\tilde{\jmath}$ contains a negative term for $\tilde{\jmath} = \tilde{\imath}$ and positive terms for $\tilde{\jmath}\neq\tilde{\imath}$, and then using Eqs. (4,5) to show that the sum over $\tilde{\jmath}$ is not positive. This is not the case if $\widetilde{n} < \widetilde{m}$, because not all sums over $\tilde{\jmath}$ contain a term with $\tilde{\jmath} = \tilde{\imath}$. However for $\widetilde{n} < \widetilde{m}$ one can do the $\tilde{\imath}$ sum first; using the same logic as above to find that the sum over $\tilde{\imath}$ is negative (or zero) for all $\tilde{\jmath}$. Thus $\sum_{\tilde{\imath}=\widetilde{n}}^{2K_{\rm n}}\sum_{\tilde{\jmath}=\widetilde{m}}^{2K_{\rm n}}\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\leq 0$ for all $\widetilde{n},\widetilde{m}$. Combining this with the fact that $\Delta\xi^{(\widetilde{n})}\geq 0$ for all $\widetilde{n}$ and $\Delta f^{(\widetilde{m})}\leq 0$ for all $\widetilde{m}$, tells us that $Y(\epsilon)\geq 0$, and thus the second law, Eq. (2), is obeyed.

As an aside, I mention that the above proof that $\sum_{\tilde{\imath}=\widetilde{n}}^{2K_{\rm n}}\sum_{\tilde{\jmath}=\widetilde{m}}^{2K_{\rm n}}\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}}(\epsilon) - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]\leq 0$ is most easily seen graphically. Write out the matrix $\left[\mathcal{T}_{\tilde{\imath}\tilde{\jmath}} - N_i\delta_{\tilde{\imath}\tilde{\jmath}}\right]$ explicitly and draw a rectangle around the elements summed over for given $\widetilde{n},\widetilde{m}$. One sees that for $\widetilde{n} > \widetilde{m}$, every *row* of the rectangle contains a diagonal ($\tilde{\imath} = \tilde{\jmath}$) element, so Eqs. (4,5) tell us that summing the elements in each *row* gives a negative (or zero) result. In contrast, ones sees that for $\widetilde{n} < \widetilde{m}$, every *column* of the rectangle contains a diagonal ($i = j$) element, thus summing over the elements in each *column* gives a negative (or zero) result.

C. Positivity of Joule heating

A very common situation is that where all reservoirs have the same temperature, so $T_i = T_0$ for all i. Then the second law, Eq. (2), reduces to the statement that the Joule heating radiated by any system, Eq. (11), will obey $J_{\rm total}\geq 0$.

VII. QUANTUM BOUND ON HEAT-CURRENTS

Here I will take Eq. (7) for the heat-current from the quantum system into reservoir i, and show that there is a negative lower-bound on its value. This shows that there is an upper-bound on the amount of heat that the DC thermoelectric response of a quantum system can suck from any given reservoir. This is irrespective of both the nature of the quantum system, and how much power one supplies to it.

The proof relies upon splitting Eq. (7) into two parts, with $J_i^{(\rm a)}$ containing only those parts of the integrals where $(\epsilon - \mu e^{-}V_i) > 0$, and $J_i^{(\rm b)}$ containing only those parts of the integrals where $(\epsilon - \mu e^{-}V_i) < 0$. In what follows I provide the proof for positive $e^{-}V_i$, however the proof for negative $e^{-}V_i$ follows exactly the same logic (under the interchange e $\leftrightarrow$ h). For positive $e^{-}V_i$, one has

$J_i = J_i^{(\mathrm{a})} + J_i^{(\mathrm{b})}$ where

$$J_i^{(\mathrm{a})}(e^{\text{-}}V_i > 0) = \int_{e^{\text{-}}V_i}^{\infty} \frac{\mathrm{d}\epsilon}{h} (\epsilon - e^{\text{-}}V_i) \sum_{j,\nu} \left[\mathcal{T}_{ij}^{+1,\nu}(\epsilon) - N_i \delta_{ij}^{+1,\nu}\right] f_j^{\nu}(\epsilon) + \int_{0}^{\infty} \frac{\mathrm{d}\epsilon}{h} (\epsilon + e^{\text{-}}V_i) \sum_{j,\nu} \left[\mathcal{T}_{ij}^{-1,\nu}(\epsilon) - N_i \delta_{ij}^{-1,\nu}\right] f_j^{\nu}(\epsilon),$$

$$J_i^{(\mathrm{b})}(e^{\text{-}}V_i > 0) = \int_{0}^{e^{\text{-}}V_i} \frac{\mathrm{d}\epsilon}{h} (\epsilon - e^{\text{-}}V_i) \sum_{j,\nu} \left[\mathcal{T}_{ij}^{+1,\nu}(\epsilon) - N_i \delta_{ij}^{+1,\nu}\right] f_j^{\nu}(\epsilon).$$

where the ± 1 is for e and h respectively. Our objective is to find the lower-bound on $J_i^{(\mathrm{a})}$ and $J_i^{(\mathrm{b})}$.

For $J_i^{(\mathrm{a})}(e^{\text{-}}V_i > 0)$, the first term in both integrands (the terms of the form $(\epsilon \pm e^{\text{-}}V_i)$) is positive, while Eq. (4) tells us that $\mathcal{T}_{ij}^{\mu\nu} > 0$, and the Fermi function satisfies $0 \leq f_j^{\nu}(\epsilon) \leq 1$. Thus $J_i^{(\mathrm{b})}(e^{\text{-}}V_i > 0)$ can never be more negative than if $f_j^{\nu}(\epsilon) = 0$ for all $j \neq i$ and $\nu \neq \mu$ (where $\mu = +1$ in the first integral and $\mu = -1$ in the second). The remaining terms — those with $j = i$ and $\mu = \nu$ — can never be more negative than if $\mathcal{T}_{ii}^{\mu\mu} = 0$. Making the substitution $\xi = (\epsilon - \mu e^{\text{-}}V_i)/(k_{\mathrm{B}}T_i)$ for $\mu = +1$ in the first integral and -1 in the second, one arrives at

$$J_i^{(\mathrm{a})}(e^{\text{-}}V_i > 0) \geq -\frac{N_i (k_{\mathrm{B}}T_i)^2}{h} \left[P(0) + P\left(\frac{e^{\text{-}}V_i}{k_{\mathrm{B}}T_i}\right)\right], \quad (16)$$

having defined $P(x) \equiv \int_x^{\infty} \mathrm{d}\xi \; \xi/(1 + \exp[\xi])$.

For $J_i^{(\mathrm{b})}(e^{\text{-}}V_i > 0)$, the first term in the integrand (the term of the form $(\epsilon - e^{\text{-}}V_i)$) is negative. Thus $J_i^{(\mathrm{b})}(e^{\text{-}}V_i > 0)$ is most negative when the j,ν-sum is as positive as possible, and this is when the Fermi functions $f_j^{\nu}(\epsilon) = 1$ for all terms with $j \neq i$ and $\nu \neq +1$. Then using Eq. (5) to write the sum over $(j,\nu) \neq (i,\mu)$ in terms of the (i,μ) element of the sum, one sees that $J_i^{(\mathrm{b})} \geq \int_0^{e^{\text{-}}V_i} \frac{\mathrm{d}\epsilon}{h} (\epsilon - e^{\text{-}}V_i)\left[N_i - \mathcal{T}_{ii}^{+1,+1}(\epsilon)\right] \left[1 - f_i^{+1}(\epsilon)\right]$, which is most negative when $\mathcal{T}_{ii}^{+1,+1} = 0$. Taking this lower bound, one can make the substitution $\xi = (\epsilon - e^{\text{-}}V_i)/(k_{\mathrm{B}}T_i)$, and use the fact that $[1 - f(\xi)] = f(-\xi)$, thus arriving at

$$J_i^{(\mathrm{b})}(e^{\text{-}}V_i > 0) \geq -\frac{N_i (k_{\mathrm{B}}T_i)^2}{h} \left[P(0) - P\left(\frac{e^{\text{-}}V_i}{k_{\mathrm{B}}T_i}\right)\right], \quad (17)$$

Taking the sum of Eqs. (16,17), and noting that $P(0) = -\mathrm{Li}_2(-1) = \pi^2/12$, where $\mathrm{Li}_2(z)$ is a dilogarithm function, one arrives at Eq. (3). Redoing the derivation for $e^{\text{-}}V_i < 0$, gives different results for $J_i^{(a)}$ and $J_i^{(b)}$, but their sum is the same as for $e^{\text{-}}V_i > 0$, thus the inequality in Eq. (3) holds for all V_i.

VIII. PHENOMENOLOGICAL TREATMENT OF DECOHERENCE AND RELAXATION

The scattering theory used above includes electrostatic interaction effects, but not dynamic interaction effects. This means that neither temperature nor bias induce the decoherence or relaxation (thermalization) of the electrons as they pass through the quantum system. However, one can include these effects in the model in a *phenomenological* manner. The idea is to model a system with such dynamic interaction effects as follows.

- Relaxation to a thermal state (and the associated decoherence) can be modeled by fictitious reservoirs[25,47,48], whose bias and temperature are chosen to ensure that the average charge and heat currents into each such reservoir is zero.
- "Pure-dephasing" (decoherence without any energy relaxation) can be modeled by an infinite set of fictitious reservoirs coupled to the system via leads which only lets a given energy pass through it, and tuning the bias on those reservoirs so that the charge and heat current into all of them are zero. The energy filtering in these leads can be treated as part of the system's scattering matrix.

A system without dynamic interactions, but with the fictitious reservoirs, obeys the bounds in Eqs. (1-3). Since the average charge and heat currents into the fictitious reservoirs are zero, the bounds on flows into the real reservoirs are the same as in the absence of the fictitious reservoirs. Thus it seems reasonable to conclude that Eqs. (1-3), are obeyed regardless of whether the electrons decohere and relax within the quantum system (due to electron-electron interactions) or not.

IX. LIMITS ON QUANTUM HEAT-ENGINES

Consider a two terminal thermoelectric device used to extract electrical power P_{total} from a heat flow between a hot reservoir (temperature T_{H}) and a cold reservoir (temperature T_{C}). The heat-current into the hot and cold reservoirs are $J_{\mathrm{H}} < 0$ and $J_{\mathrm{C}} > 0$, while the power generated, $P_{\mathrm{total}} > 0$. Combining the first and second laws, Eqs. (1,2), one finds Carnot's result that

$$P_{\mathrm{total}} \leq -J_{\mathrm{H}} \left(1 - T_{\mathrm{C}}/T_{\mathrm{H}}\right). \quad (18)$$

This inequality limits the efficiency, $\eta = P_{\mathrm{total}}/(-J_{\mathrm{H}})$, with which the heat-engine extracts electrical power from a given heat-current. The equality corresponds to Carnot efficiency. Combining Eq. (18) with the bound on J_{H} in Eq. (3), one gets the absolute limit on the power

$$P_{\mathrm{total}} \leq \frac{N_{\mathrm{H}} \pi^2 k_{\mathrm{B}}^2 T_{\mathrm{H}} (T_{\mathrm{H}} - T_{\mathrm{C}})}{6h}. \quad (19)$$

For a heat-engine between a hot-source at $T_{\mathrm{H}} = 100°\mathrm{C} = 373\mathrm{K}$, and a cold source at $T_{\mathrm{C}} = 20°\mathrm{C} = 293\mathrm{K}$, the maximum power per transverse mode is 15nW.

X. LIMITS ON QUANTUM REFRIGERATORS

Consider a two terminal device extracting heat from a cold reservoir (to keep it cold or cool it further), when the ambient temperature is T_0 and the cold reservoir's temperature is $T_{\rm C} < T_0$. Power is supplied (negative $P_{\rm total}$) to extract heat from the cold reservoir (negative $J_{\rm C}$). Combining the first and second laws, Eqs. (1,2), one gets the Carnot bound on refrigeration. Eq. (3) gives a second quantum bound. Together they read,

$$-J_{\rm C} \leq \begin{cases} -P_{\rm total}\, T_{\rm C}/(T_0 - T_{\rm C}), & \text{[thermodyn.]} \\ N_{\rm C}\pi^2 (k_{\rm B}T_{\rm C})^2/6h. & \text{[quantum]} \end{cases} \tag{20}$$

The thermodynamic bound places no limit on the heat that one can extract from the cold reservoir; one can always extract heat if one is able to provide enough electrical power. In stark contrast the quantum bound places an absolute limit on the heat extracted.

As a first example, consider a kitchen refrigerator running at 100W, cooling from room-temperature $T_0 = 293$K to $T_{\rm C} = 0°$C$= 273$K, and now imagine it consists of thermoelectric devices. Then the bounds are

$$-J_{\rm C} \leq \begin{cases} 1300\,{\rm W}, & \text{[thermodyn.]} \\ N_{\rm C} \times 36\,{\rm nW}. & \text{[quantum]} \end{cases} \tag{21}$$

The quantum bound is stronger than the thermodynamic one if $N_{\rm C} \lesssim 10^{10}$. A typical semiconductor thermoelectric (Fermi wavelength ~ 100nm) with a cross-section of 1cm×1cm, has $N_{\rm C} \sim 10^{10}$; the quantum bound dominates for cross-sections smaller than this.

As a second example, consider a point-contact nanostructure such as in Ref. [54]. There typical biases and currents are tens of mV and tenths of nA, so typical powers are pico-Watts (pW). Suppose such a device is used for cooling as proposed in Ref. [13], to refrigerate a micron-sized island from the cryostat temperature $T_0 \sim 1$K down to $T_{\rm C} \sim 0.1$K. Then

$$-J_{\rm C} \leq \begin{cases} 0.1\,{\rm pW}, & \text{[thermodyn.]} \\ N_{\rm C} \times 0.005\,{\rm pW}. & \text{[quantum]} \end{cases} \tag{22}$$

For a single-mode point contact $N_{\rm C} \sim 1$ it is clear that the thermodynamic bound is irrelevant. However it becomes relevant if one has 20 or more such single-mode point-contacts in parallel thermally (for example as sketched in Fig. 2b of Ref. [13]).

XI. NERNST'S UNATTAINABILITY PRINCIPLE

Nernst's unattainability principle, is one of the two statements which are together referred to as the third law of thermodynamics. It says that no matter how much work one does, one cannot take a reservoir initially at finite temperature to zero temperature in a finite time (the other part of the third law is the claim that entropy vanishes at zero temperature). In the context of a reservoir i which is a gas of free electrons in equilibrium, we use the textbook result that the heat capacity of a such a Fermi gas, $C_i \equiv ({\rm d}Q_i/{\rm d}T_i) \propto T_i$. If the quantum system is sucking heat out over a long time via a DC bias, then the fastest rate of heat removal is given by quantum bound, $J_i^{\rm qb} \propto T_i^2$. Assuming this heat flow is weak enough that the reservoir remains in equilibrium, we have $-J_i^{\rm qb} = ({\rm d}Q_i/{\rm d}t) = C_i({\rm d}T_i/{\rm d}t)$. Thus

$$\frac{{\rm d}T_i}{{\rm d}t} = -T_i^{\zeta}, \qquad \text{with } \zeta = 1. \tag{23}$$

Solving this for arbitrary ζ, one sees that $T_i = 0$ is reached in a finite time if $\zeta < 1$, but not if $\zeta > 1$. Eq. (23) has $\zeta = 1$, which is the critical value of ζ, then the temperature decays exponentially with time t; thus it never reach absolute zero, but does get exponentially close to it. A concrete example with $\zeta = 1$ is the point-contact considered in Ref. [13] at large bias. For $\zeta > 1$ (e.g. the pump in Ref. [29] has $\zeta = 3/2$ for a fermionic bath) the decay is much slower, going like $t^{-1/(\zeta-1)}$ for large t.

Ref. [30] proposes a quantum system with $\zeta < 1$, violating the unattainability principle. It is clear that their system does not fall into the broad class of systems considered here. In particular the bosonic reservoir they cool is very different from the ones that I consider.

XII. CONCLUDING REMARKS

This work is on the DC response of any quantum device where single-electron charging effects are not significant (single electron charging energy much less than temperature or level-broadening due to the coupling to reservoirs). Here "DC" means the zero frequency response to time-independent potentials. However, many classical heat-engines and refrigerators rely on a pumping cycle, so it would be worthwhile to extend this analysis to the quantum equivalents of such finite-frequency driven systems (quantum pumps, etc). One should also consider that quantum transport is inherently noisy, so the energy and entropy in the quantum system fluctuates randomly on short timescales. These fluctuations give little contribution to low-frequency responses, because they average out on long timescales. In the future we hope to address whether such quantum fluctuations obey the classical fluctuation theorem[55].

Finally, it is worth recalling that all the results presented here rely on only two ingredients; (i) Schrodinger's equation formulated as a scattering theory, and (ii) simple *equilibrium* statistical mechanics in the reservoirs. It is remarkable that the thermodynamic laws for *non-equilibrium* transport (far outside the linear-response regime) emerge so naturally from a simple combination of these two ingredients.

[1] H.J. Goldsmid, *Introduction to Thermoelectricity* (Springer, Heidelberg, 2009).

[2] F.J. DiSalvo, Science **285**, 703 (1999).

[3] A. Shakouri and M. Zebarjadi Chapt 9 of *Thermal nanosystems and nanomaterials*, S. Volz (Ed.) (Springer, Heidelberg, 2009). A. Shakouri , Annu. Rev. Mater. Res. **41**, 399 (2011).

[4] F. Giazotto, T.T. Heikkila, A. Luukanen, A.M. Savin, and J.P. Pekola, Rev. Mod. Phys. **78**, 217 (2006). J.T. Muhonen, M. Meschke, and J.P. Pekola, Rep. Prog. Phys. **75**, 046501 (2012).

[5] M. Zebarjadi, K. Esfarjani, and A. Shakouri, Appl. Phys. Lett. **91**, 122104 (2007); Mater. Res. Soc. Proc. **1044** U10-04 (2008).

[6] M. Wierzbicki and R. Swirkowicz, Phys. Rev. B **84**, 075410 (2011).

[7] T. Gunst, T. Markussen, A.-P. Jauho, and M. Brandbyge, Phys. Rev. B **84**, 155449 (2011).

[8] G. Rajput and K.C. Sharma, J. Appl. Phys. **110**, 113723 (2011).

[9] A. Glatz, N.M. Chtchelkatchev, I.S. Beloborodov, and V. Vinokur, Phys. Rev. B **84**, 235101 (2011).

[10] B. Sothmann, R. Sánchez, A.N. Jordan, and M. Büttiker, Phys. Rev. B 85, 205301 (2012).

[11] B. Sothmann and M. Büttiker, EPL 99, 27001 (2012).

[12] P. Trocha and J. Barnaś, Phys. Rev. B **85**, 085408 (2012).

[13] R.S. Whitney, preprint arXiv:1208.6130.

[14] G. Casati, C. Mejía-Monasterio, and T. Prosen, Phys. Rev. Lett. **101**, 016601 (2008).

[15] D. Nozaki, H. Sevinçli, W. Li, R. Gutiérrez, and G. Cuniberti, Phys. Rev. B **81**, 235406 (2010).

[16] O. Entin-Wohlman, Y. Imry, and A. Aharony, Phys. Rev. B **82**, 115314 (2010).

[17] K.K. Saha, T. Markussen, K.S. Thygesen, and B.K. Nikolić, Phys. Rev. B **84**, 041412(R) (2011).

[18] O. Karlström, H. Linke, G. Karlström, and A. Wacker, Phys. Rev. B **84**, 113415 (2011).

[19] Y.-S. Liu, B.C. Hsu, and Y.-C. Chen, J. Phys. Chem. C **115**, 6111 (2011).

[20] J. Gemmer, M. Michel, and G Mahler, Quantum Thermodynamics (Springer-Verlag, Berlin, 2010).

[21] J.C. Wheeler, Phys. Rev. A **43**, 5289 (1991).

[22] S. Abe and A.K. Rajagopal, Phys. Rev. Lett. **91**, 120601 (2003).

[23] F.G.S.L. Brandão and M.B. Plenio, Nature Phys. **4**, 873 (2008).

[24] T. Sagawa and M. Ueda, Phys. Rev. Lett. **100**, 080403 (2008). K. Jacobs, Phys. Rev. A **80**, 012322 (2009).

[25] Ph.A. Jacquet, J. Stat. Phys. **134**, 709 (2009).

[26] L. del Rio, J. Aberg, R. Renner, O. Dahlsten, and V. Vedral, Nature **474**, 61 (2011).

[27] L. Bruneau, V. Jakšić, and C.-A. Pillet, preprint arXiv:1201.3190.

[28] K. Maruyama, F. Nori, and V. Vedral, Rev. Mod. Phys. **81**, 1 (2009).

[29] A. Levy, R. Alicki, and R. Kosloff, Phys. Rev. E **85**, 061126 (2012).

[30] M. Kolar, D. Gelbwaser-Klimovsky, R. Alicki, and G. Kurizki, Phys. Rev. Lett. **109**, 090601 (2012).

[31] In this work "DC" is used to mean the zero-frequency heat and charge currents due to time-independent biases and temperature differences. In Sect. XII, I briefly mention the limitations of studying only this DC response.

[32] Ref. [25] discusses the second law within scattering theory, but only when $I \propto V$ with no thermoelectric effects.

[33] S. Datta, *Electronic Transport in Mesoscopic Systems* (Cambridge University Press, Cambridge, 1997).

[34] Ya.M. Blanter and M. Büttiker, Phys. Rep. **336**, 1 (2000).

[35] M. Büttiker, J. Phys. Condens. Matter, **5**, 9361 (1993). M. Büttiker, A. Prêtre, and H. Thomas, Phys. Rev. Lett. **70**, 4114 (1993); Z. Phys. B, **94**, 133 (1994). T. Christen, and M. Büttiker, Phys. Rev. Lett. **77**, 143 (1996).

[36] C. Petitjean, D. Waltner, J. Kuipers, I. Adagideli, K. Richter, Phys. Rev. B, **80**,115310 (2009).

[37] T. Christen and M. Büttiker, Europhys. Lett. **35**, 523 (1996).

[38] D. Sanchez and M. Buttiker, Phys. Rev. Lett. **93**, 106802 (2004).

[39] J. Meair and Ph. Jacquod, J. Phys.: Cond. Matter **24**, 272201 (2012).

[40] D. Sanchez and R. Lopez, preprint arXiv:1209.1264.

[41] H.-L. Engquist and P.W. Anderson, Phys. Rev. B **24**, 1151 (1981).

[42] U. Sivan and Y. Imry, Phys. Rev. B **33**, 551 (1986).

[43] P.N. Butcher, J. Phys.: Condens. Matter, **2**, 4869 (1990). His Eq. (16c) contains "$\mathrm{d}\varepsilon_\beta$", it should be "$\mathrm{d}\varepsilon\, f_\beta$".

[44] N.R. Claughton and C.J. Lambert, Phys. Rev. B **53**, 6605 (1996).

[45] Ph. Jacquod and R.S. Whitney, Europhys. Lett. **91**, 67009 (2010).

[46] Ph. Jacquod, R.S. Whitney, Jonathan Meair, and M. Büttiker, Phys. Rev. B **86**, 155118 (2012).

[47] M. Büttiker, Phys. Rev B **33**, 3020 (1986); IBM J. Res. Dev. **32**, 63 (1988).

[48] Ph. A. Jacquet and C.-A. Pillet, Phys. Rev. B **85**, 125120 (2012).

[49] The Stefan-Boltzmann law for *relativistic* fermions (neutrinos)[56] at $k_{\mathrm{B}}T \gg E_{\mathrm{F}}$ has $J \propto T^4$. All these Stefan-Boltzmann laws take the form $J \propto N\ (k_{\mathrm{B}}T)^2$, with $N \sim A/\lambda_{\mathrm{deBr}}^2$ for aperture cross-section A. For non-relativistic fermions with $E_{\mathrm{F}} \gg k_{\mathrm{B}}T$, the de Broglie wavelength is $\lambda_{\mathrm{deBr}} \sim h/\sqrt{2mE_{\mathrm{F}}}$. In contrast, relativistic fermions with $k_{\mathrm{B}}T \gg E_{\mathrm{F}}$ and photons both have $\lambda_{\mathrm{deBr}} \sim hc/(k_{\mathrm{B}}T)$.

[50] F. Belgiorno, J. Phys. A: Math. Gen. **36**, 8165 (2003).

[51] P.T. Landsberg, Rev. Mod. Phys. **28**, 363 (1956).

[52] I assume the sc reservoir is a superconducting condensate without quasi-particles, i.e. sc gap much larger than temperatures and biases. Quasiparticles in the superconductor can be treated as a separate electronic reservoir.

[53] Upon measuring the energy ϵ relative to $e^{-}V_{\mathrm{sc}}$, the condition for gauge-invariance[37] becomes $\sum_{k'} \left(\mathrm{d}\mathcal{T}_{ij}^{\mu\nu}(\epsilon, \mathbf{v}_{\mathrm{env}})/\mathrm{d}V_{k'}\right) = 0$ for all ϵ and $\mathbf{v}_{\mathrm{env}}$, where the sum is over all reservoir and gate voltages (including V_{sc}). This is satisfied by taking all potentials (including the screening potentials) as relative to V_{sc}.

[54] L.P. Kouwenhoven, B.J. van Wees, C.J.P.M. Harmans, J.G. Williamson, H. van Houten, C.W.J. Beenakker, C.T. Foxon, and J.J. Harris, Phys. Rev. B **39**, 8040 (1989).

[55] D.J. Evans and D.J. Searles, Phys. Rev. E **50**, 1645 (1994). S.K. Blau, Phys. Today **55**, 19 (2002).

[56] see e.g. B. Kuchowicz, Lett. Nuovo Cimento **8**, 942 (1973).